

Dirac's Constrained Hamiltonian Dynamics from an Unconstrained Dynamics

Heinz J. Rothe

Institut für Theoretische Physik

Universität Heidelberg, Philosophenweg 16, D-69120 Heidelberg, Germany ¹

February 1, 2008

Abstract

We derive the Hamilton equations of motion for a constrained system in the form given by Dirac, by a limiting procedure, starting from the Lagrangean for an unconstrained system. We thereby elucidate the role played by the primary constraints and their persistence in time.

The Hamiltonian formulation for systems whose dynamics is described by a Lagrangean with singular Hessian has been given a long time ago by Dirac [1], and has been elaborated ever since in numerous papers ². As is well known all gauge theories fall into the class of singular systems. The usual starting point for deriving the Hamilton equations of motion is the singular Lagrangean which leads, in the language of Dirac, to primary constraints. While the primary constraints have no analog on the Lagrangean level, requiring their persistence in time leads to equations relating the coordinates and velocities. These are part of the Euler-Lagrange equations of motion. This persistence requirement may lead to secondary constraints and is the first step in the Dirac algorithm generating the constraints. The purpose of

¹email: h.rothe@thphys.uni-heidelberg.de

²It is impossible to quote here the very large number of papers. Comprehensive discussions can be found e.g. in [2]

this paper is to derive the Hamilton equations of motion for systems with a singular Hessian, by starting from an unconstrained system and taking an appropriate limit. The formulation of the equations of motion within an extended phase space is quite natural, and the role played by the primary constraints, and the demand for their persistence in time will be illuminated thereby.

Consider the following Lagrangean quadratic in the velocities

$$L = \frac{1}{2} \sum_{ij} W_{ij}(q; \alpha) \dot{q}_i \dot{q}_j - \sum_i \eta_i(q) \dot{q}_i - V(q) . \quad (1)$$

where q stands for the set of coordinates $\{q_i\}$ and α stands collectively for a set of parameters. For $\alpha \neq \alpha_c$ we assume that $\det W \neq 0$, so that we are dealing with a non-singular system. We assume that the singular system of interest is realized for $\alpha = \alpha_c$ (the subscript c stands for "critical") where $\det W(q; \alpha_c) = 0$. Our aim is to derive the Hamilton equations of motion for the singular system, by taking the limit $\alpha \rightarrow \alpha_c$ of the Hamilton equations of motion following from (1).³ For $\alpha \neq \alpha_c$, one readily constructs the Hamiltonian

$$H = \frac{1}{2} \sum_{ij} (p_i + \eta_i) W_{ij}^{-1} (p_j + \eta_j) + V(q) , \quad (2)$$

where the canonical momenta are related to the velocities by

$$p_i = \sum_j W_{ij}(q; \alpha) \dot{q}_j - \eta_i(q) . \quad (3)$$

The symmetric matrix W can be diagonalized by an orthogonal transformation, $W_D = C^T W C$ with $C_{i\ell} = v_i^{(\ell)}$, where $\vec{v}^{(\ell)}$ are the orthonormalized eigenvectors of W ,

$$W(q; \alpha) \vec{v}^{(\ell)}(q; \alpha) = \lambda_\ell(q; \alpha) \vec{v}^{(\ell)}(q; \alpha) . \quad (4)$$

In terms of the eigenvalues and eigenvectors, W can be written in the form

$$W_{ij} = \sum_\ell \lambda_\ell v_i^{(\ell)} v_j^{(\ell)} . \quad (5)$$

³Clearly the form of (1) is not unique.

Correspondingly W_{ij}^{-1} is obtained by making the replacement $\lambda_\ell \rightarrow \frac{1}{\lambda_\ell}$. Hence the Hamiltonian (2) is given by

$$H = \frac{1}{2} \sum_\ell \frac{1}{\lambda_\ell} \phi_\ell^2 + V(q) , \quad (6)$$

where

$$\phi_\ell(q, p; \alpha) := (\vec{p} + \vec{\eta}(q)) \cdot \vec{v}^{(\ell)}(q; \alpha) . \quad (7)$$

For $\alpha \neq \alpha_c$ the Hamilton equations of motion then take the form

$$\dot{q}_i = \frac{\partial H}{\partial p_i} = \sum_\ell \frac{1}{\lambda_\ell} \phi_\ell \frac{\partial \phi_\ell}{\partial p_i} , \quad (8)$$

$$\dot{p}_i = -\frac{\partial H}{\partial q_i} = -\frac{1}{2} \sum_\ell \frac{\partial}{\partial q_i} \left(\frac{1}{\lambda_\ell} \phi_\ell^2 \right) - \frac{\partial V}{\partial q_i} . \quad (9)$$

Consider now the limit $\alpha \rightarrow \alpha_c$ where $\det W = 0$. Let $\{\lambda_{\ell_0}\}$ denote the set of eigenvalues which vanish in this limit. In order to implement the limit we first write (8) and (9) in the form

$$\dot{q}_i = \sum_{\ell \neq \{\ell_0\}} \frac{1}{\lambda_\ell} \phi_\ell \frac{\partial \phi_\ell}{\partial p_i} + \sum_{\ell_0} \frac{1}{\lambda_{\ell_0}} \phi_{\ell_0} \frac{\partial \phi_{\ell_0}}{\partial p_i} , \quad (10)$$

$$\dot{p}_i = -\frac{1}{2} \sum_{\ell \neq \{\ell_0\}} \frac{\partial}{\partial q_i} \left(\frac{1}{\lambda_\ell} \phi_\ell^2 \right) - \frac{1}{2} \sum_{\ell_0} \frac{\partial}{\partial q_i} \left(\frac{1}{\lambda_{\ell_0}} \phi_{\ell_0}^2 \right) - \frac{\partial V}{\partial q_i} . \quad (11)$$

Now, from (3) and (7) we have that

$$\phi_{\ell_0} = \sum_i v_i^{(\ell_0)} W_{ij} \dot{q}_j = \lambda_{\ell_0} \vec{v}^{(\ell_0)} \cdot \dot{\vec{q}} . \quad (12)$$

Hence

$$\lim_{\alpha \rightarrow \alpha_c} \frac{1}{\lambda_{\ell_0}} \phi_{\ell_0} = \rho_{\ell_0}(q, \dot{q}; \alpha_c) , \quad (13)$$

where

$$\rho_{\ell_0}(q, \dot{q}; \alpha_c) = \dot{\vec{q}} \cdot \vec{v}^{(\ell_0)}(q; \alpha_c) . \quad (14)$$

Note that the finiteness of the velocities in (12) implies that

$$\phi_{\ell_0}(q, p, \alpha_c) = 0 , \quad (15)$$

which are just the primary constraints. Hence for $\alpha \rightarrow \alpha_c$ (10) reduces to

$$\dot{q}_i = \sum_{\ell \neq \{\ell_0\}} \left(\frac{1}{\lambda_\ell} \phi_\ell \frac{\partial \phi_\ell}{\partial p_i} \right)_{\alpha=\alpha_c} + \sum_{\ell_0} \rho_{\ell_0} \left(\frac{\partial \phi_{\ell_0}}{\partial p_i} \right)_{\alpha=\alpha_c} . \quad (16)$$

Consider next eqs. (11). Making again use of (12) and the fact that $\frac{\partial \lambda_{\ell_0}}{\partial q_i} |_{\alpha=\alpha_c} = 0$ one obtains that for $\alpha \rightarrow \alpha_c$ these reduce to

$$\dot{p}_i = -\frac{1}{2} \sum_{\ell \neq \{\ell_0\}} \left[\frac{\partial}{\partial q_i} \left(\frac{1}{\lambda_\ell} \phi_\ell^2 \right) \right]_{\alpha=\alpha_c} - \sum_{\ell_0} \rho_{\ell_0} \left(\frac{\partial \phi_{\ell_0}}{\partial q_i} \right)_{\alpha=\alpha_c} - \frac{\partial V}{\partial q_i} , \quad (17)$$

where use has been made of (13).

We now notice that the first sum on the RHS of eqs. (16) and (17) are just given by $\frac{\partial H_c}{\partial p_i}$ and $-\frac{\partial H_c}{\partial q_i}$, where H_c is the canonical Hamiltonian obtained from (6) by taking the limit $\alpha \rightarrow \alpha_c$:

$$H_C = \frac{1}{2} \sum_{\ell \neq \ell_0} \frac{1}{\lambda_\ell(q, \alpha_c)} \phi_\ell^2(q, p; \alpha_c) + V(q) \quad (18)$$

Here use has again be made of (15) and of the fact that $\lim_{\alpha \rightarrow \alpha_c} \frac{\phi_{\ell_0}}{\lambda_{\ell_0}}$ is finite. H_C is just the canonical Hamiltonian derived from the (1) for $\alpha = \alpha_c$, evaluated on the primary surface. Hence the equations of motion (16) and (17) take the well known form

$$\dot{q}_i = \frac{\partial H_T}{\partial p_i} \quad (19)$$

$$\dot{p}_i = -\frac{\partial H_T}{\partial q_i} \quad (20)$$

where H_T is the "total" Hamiltonian

$$H_T = H_C + \sum_{\ell_0} \rho_{\ell_0} \phi_{\ell_0} , \quad (21)$$

and ρ_{ℓ_0} are the undetermined projections of the velocities on the zero modes (14). Note that the derivatives in (19) and (20) are understood not to act on ρ_{ℓ_0} . Eqs. (19) and (20) must be supplemented by the primary constraints

(15). In fact these equations only have a solution if $q_i(t)$ and $p_i(t)$ are points in the submanifold defined by the primary constraints [3]. Note that from (16) it follows that the projection of $\dot{\vec{q}}$ onto the zero mode $\vec{v}^{(\ell_0)}$ reduces to an identity, since $\sum_i v_i^{(\ell_0)} \frac{\partial H_c}{\partial p_i} = 0$. Hence we have obtained the Hamilton equations of motion for a constrained system in the form given by Dirac, by taking the limit $\alpha \rightarrow \alpha_c$ of the equations of motion for an unconstrained system. The primary constraints are just the statement that the projected velocities (14) are finite in this limit.

Actually, eqs. (19) and (20), together with the primary constraints (15) do not directly yield the complete set of Lagrange equations of motion. These follow by also implementing the persistence in time of the primary constraints. The primary constraints themselves have no analog on the Lagrangean level but allow us to recover the connection between momenta and velocities needed to express the Hamilton equations of motion in terms of Lagrangean variables. Thus from (16) and (7) it follows that

$$\dot{q}_i = \sum_{\ell \neq \{\ell_0\}} \frac{1}{\lambda_\ell} \phi_\ell v_i^{(\ell)} + \sum_{\ell_0} (\dot{\vec{q}} \cdot v^{(\ell_0)}) v_i^{(\ell_0)} , \quad (22)$$

where it is understood that we have set $\alpha = \alpha_c$. Define the matrix W_{ij} constructed from $v^{(\ell)}$ and λ_ℓ ($\ell \neq \{\ell_0\}$): $W_{ij} = \sum_{\ell \neq \{\ell_0\}} \lambda_\ell v_i^{(\ell)} v_j^{(\ell)}$. From (22) we obtain

$$\sum_j W_{ij}(q, \alpha_c) \dot{q}_j = \sum_{\ell, j} (p + \eta)_j v_i^{(\ell)} v_j^{(\ell)} = (p + \eta)_i , \quad (23)$$

where we have made use of the primary constraints (15), in order to extend the sum on the RHS of (23) over all ℓ , and of the completeness relation for the eigenvectors. Hence we have recovered (3) for $\alpha = \alpha_c$.

As we have pointed out above, the persistence of the primary constraints yields on Lagrange level equations involving only coordinates and velocities. These are part of the Euler-Lagrange equations of motion. From the point of view taken in this paper, that the equations of motion are obtained by a limiting procedure from those of an unconstrained system, the persistence of

the primary constraints can also be viewed to follow from the requirement that also the accelerations remain finite in the limit $\alpha \rightarrow \alpha_c$. Thus for $\alpha \neq \alpha_c$ we have from (12) that $\phi_{\ell_0} = \lambda_{\ell_0} \dot{\vec{q}} \cdot \vec{v}^{(\ell_0)}$. Taking the time derivative of this expression, and noting that $\lim_{\alpha \rightarrow \alpha_c} \lambda_{\ell_0}(q, \alpha) = 0$, and $\lim_{\alpha \rightarrow \alpha_c} \partial_i \lambda_{\ell_0}(q, \alpha) = 0$, we immediately conclude that $\dot{\phi}_{\ell_0}(q, p, \alpha_c) = 0$. On the Hamiltonian level this requirement must be implemented explicitly to yield the missing Euler-Lagrange equations of motion, not manifest in (19) and (20).

From the above discussion it is evident that the limit $\alpha \rightarrow \alpha_c$ must be carried out on the level of the Hamilton *equations of motion* of the unconstrained system, whereas on the Lagrangean level we are allowed to take this limit directly in the Lagrangean. The equivalence between the Lagrangean and Hamiltonian formulations has been studied in detail in [3].

As an example consider the singular Lagrangean of the pure $U(1)$ Maxwell theory:

$$L[A^\mu, \dot{A}^\mu] = -\frac{1}{4} \int d^3x F_{\mu\nu} F^{\mu\nu} . \quad (24)$$

Consider further the non-singular Lagrangean

$$L[A^\mu, \dot{A}^\mu] = -\frac{1}{4} \int d^3x F_{\mu\nu} F^{\mu\nu} + \frac{1}{2} \alpha \int d^3x \dot{A}^0 , \quad (25)$$

which for $\alpha \rightarrow 0$ reduces to (24). This (non-covariant) choice is of course only the simplest one. Any other Lagrangean reducing to (24) in the appropriate limit would be just as acceptable. The canonical momenta conjugate to A^μ are given by

$$\pi_\mu = F^{0\mu} + \alpha \delta_{\mu 0} (\partial_0 A^0) . \quad (26)$$

The Lagrangean written in the analogous form to (1) is

$$L = \int d^3x d^3y \left[\frac{1}{2} \dot{A}^\mu(\vec{x}, x^0) W_{\mu\nu}(\vec{x}, \vec{y}) \dot{A}^\nu(\vec{y}, x^0) \right] - \int d^3x \eta_\mu(A(x)) \dot{A}^\mu(x) - V[A] , \quad (27)$$

where

$$\eta_\mu := (0, -\vec{\nabla} A^0) , \quad (28)$$

and the matrix elements of the symmetrix matrix W read

$$W_{0i}(\vec{x}, \vec{y}) = 0 , \quad (29)$$

$$W_{00}(\vec{x}, \vec{y}) = \alpha \delta^{(3)}(\vec{x} - \vec{y}) , \quad (30)$$

$$W_{ij}(\vec{x}, \vec{y}) = \delta_{ij} \delta^{(3)}(\vec{x} - \vec{y}) . \quad (31)$$

The potential $V[A]$ is given by

$$V[A] = \int d^3x \left(\frac{1}{4} F_{ij} F^{ij} - \frac{1}{2} (\vec{\nabla} A^0)^2 \right) , \quad (32)$$

and the canonical momenta, analogous to (3) read

$$\pi_\mu(x) = \int d^3y W_{\mu\nu}(\vec{x}, \vec{y}) \dot{A}^\nu(\vec{y}, x^0) - \eta_\mu(A(x)) . \quad (33)$$

The matrix W possesses the following orthonormalized eigenvectors, labeled by a discrete and continuous index, replacing the discrete index ℓ in (4)

$$v_\nu^{(\rho, \vec{z})}(\vec{x}) = \delta_{\rho\nu} \delta^{(3)}(\vec{x} - \vec{z}) , \quad (34)$$

and the corresponding eigenvalues of W are given by

$$\lambda^{(0, \vec{z})} = \alpha ; \quad \lambda^{(i, \vec{z})} = 1 \quad (i = 1, 2, 3) . \quad (35)$$

The Hamiltonian, analogous to (6) then takes the form

$$H = \frac{1}{2} \sum_i \int d^3z \left[\int d^3x (\pi_\mu(x) + \eta_\mu(x)) v_\mu^{(i, \vec{z})}(\vec{x}) \right]^2 \quad (36)$$

$$+ \frac{1}{2} \int d^3z \frac{1}{\alpha} \left[\int d^3x (\pi_\mu(x) + \eta_\mu(x)) v_\mu^{(0, \vec{z})}(\vec{x}) \right]^2 + V[A] . \quad (37)$$

Upon making use of (34), this expression reduces to

$$H = \frac{1}{2} \sum_i \int d^3z (\pi_i + \eta_i)^2 + \frac{1}{2} \int d^3z \frac{1}{\alpha} (\pi_0 + \eta_0)^2 + V[A] , \quad (38)$$

where in the present case $\eta_0 = 0$. The Hamilton equations of motion read

$$\dot{A}^0 = \frac{\delta H}{\delta \pi_0} = \frac{\pi_0}{\alpha} ,$$

$$\begin{aligned}
\dot{A}^i &= \frac{\delta H}{\delta \pi_i} = \pi_i + \partial^i A^0 , \\
\dot{\pi}_0 &= -\frac{\delta H}{\delta A^0} = -\vec{\nabla} \cdot \vec{\pi} , \\
\dot{\pi}_i &= -\frac{\delta H}{\delta A^i} = -\partial_j F^{ji} .
\end{aligned} \tag{39}$$

For finite $\dot{A}^0$ the first equation tells us that in the limit $\alpha \rightarrow 0$, π_0 must vanish, whereas $\dot{A}^0$ remains completely arbitrary. Since π_0 must vanish for arbitrary times, the third equation tells us that also $\vec{\nabla} \cdot \vec{\pi} = 0$. This is just the secondary (Gauss law) constraint.

Alternatively we could have departed from a covariant form for an unconstrained system by adding to the Lagrangean density in (24) the covariant term $\frac{\alpha}{2}(\partial_\mu A^\mu)^2$. In this case $\eta_\mu(A(x))$ also depends on α , and following our general procedure one is led to the equations of motion

$$\dot{A}^0 = \frac{\partial H}{\partial \pi_0} = \frac{1}{\alpha} \pi_0 - \vec{\nabla} \cdot \vec{A} , \tag{40}$$

$$\dot{A}^i = \frac{\partial H}{\partial \pi_i} = \pi_i + \partial^i A^0 , \tag{41}$$

$$\dot{\pi}_0 = -\frac{\partial H}{\partial A^0} = -\vec{\nabla} \cdot \vec{\pi} , \tag{42}$$

$$\dot{\pi}_i = -\partial_j F^{ji} + \partial^i \pi_0 . \tag{43}$$

From the first equation it follows again that in the limit $\alpha \rightarrow 0$, $\pi_0 = 0$ for all times, so that we are left with the standard equations for the pure Maxwell theory. We emphasize once more that the limit must be taken on the level of the equations of motion, and not in the Hamiltonian. The examples demonstrate the, of course, well known fact, that only after taking into account the primary constraints and their persistence in time, the full set of Euler-Lagrange equations of motion are generated. Thus primary constraints, and possible secondary constraints following from them, play a special role, while tertiary, etc. constraints correspond to consistency relations hidden in the Euler-Lagrange equations of motion. With the primary constraints written

in the form (15), a strict iterative construction of the persistence equations for the constraints will necessarily parallel exactly the equations obtained on the Lagrangean level, irrespective of any possible bifurcations⁴.

Let us summarize. In this paper we have shown that the Hamilton equations of motion for any constrained system, in the form given by Dirac, can be obtained as a limit of the equations of motion for an unconstrained system. It was thereby shown that the primary constraints follow directly from the requirement that the velocities be finite in this limit. To obtain the full set of equations on Hamiltonian level, which translate into the Euler-Lagrange equations of motion, one must take into account the persistence of the primary constraints in time. These equations, which are implicit in the persistence requirement of the primary constraints, can also be viewed to follow from the requirement that also the accelerations remain finite in the above mentioned limit. The particular form of the term added to the singular Lagrangean which converts the system into a second class system, plays no role. The only requirement is that the unconstrained system reduces to the constrained theory of interest in an appropriate limit. We have demonstrated this for the case of the pure Maxwell theory.

Acknowledgements

I thank Klaus D. Rothe for several constructive comments.

References

- [1] P.A.M.Dirac, Can. J. Math. **2** (1950) 129; *Lectures on Quantum Mechanics* (Yeshiva University, New York, 1964).
- [2] E.C.G. Sudarshan and N. Mukunda, *Classical Dynamics: A Modern Perspective* (Wiley, New York, 1974); A. Hanson, T. Regge and C. Teit-

⁴For examples exhibiting bifurcations see [4]

elboim, *Constrained Hamiltonian Systems*, (Accademia Nazionale dei Lincei, Roma (1975)); K. Sundermeyer, *Constrained Dynamics*, Lecture Notes in Physics, Vol. **169** (Springer, Berlin (1982)); D.M. Gitman and I.V. Tyutin, *Quantization of Fields with Constraints*, Springer (1980); M. Henneaux and C. Teitelboim, *Quantization of Gauge Systems*, Princeton University Press (1992).

[3] C. Batlle, J. Gomis, J.M. Pons and N. Roman-Roy, J. Math. Phys. **27** (1986) 2953.

[4] L. Lusanna, Nuovo Cimento Vol. **14**, Nr. 3 (1991) 1